

## AMENDMENTS TO THE SPECIFICATION:

Please change the title at page 1, line 1, from "Fungicidally active compound combinations" to

-- FUNGICIDAL ACTIVE SUBSTANCE COMBINATIONS --

Please insert the following at page 1, between the title and line 3:

-- The present patent application has been filed under 35 U.S.C. 371 as a national stage application of PCT/EP2004/008072, filed July 20, 2004, which was published in German as International Patent Publication WO 2005/009130 on February 3, 2005, and is entitled to the right of priority of German Patent Application 103 33 373.8, filed July 23, 2003. --

Please insert the following at page 13, between the lines 18 and 20 (compare WO 2005/009130 at page 11, line 15, through page 13, line 12):

-- As already mentioned above, it is possible to treat all plants and their parts according to the invention. In a preferred embodiment, wild plant species and plant cultivars, or those obtained by conventional biological breeding methods, such as crossing or protoplast fusion, and parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant cultivars obtained by genetic engineering methods, if appropriate in combination with conventional methods (genetically modified organisms), and parts thereof, are treated. The term "parts" or "parts of plants" or "plant parts" has been explained above.

With particular preference, plants of the plant cultivars which are in each case commercially available or in use are treated according to the invention. Plant cultivars are to be understood as meaning plants having new properties (traits) and which have been obtained by conventional breeding, by mutagenesis or by recombinant DNA techniques. They can be cultivars, biotypes or genotypes.

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive (synergistic) effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an

increase in the activity of the substances and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products are possible which exceed the effects which were actually to be expected.

The transgenic plants or plant cultivars (i.e. those obtained by genetic engineering) which are preferably to be treated according to the invention include all plants which, in the genetic modification, received genetic material which imparted particularly advantageous useful properties (traits) to these plants. Examples of such properties are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products. Further and particularly emphasized examples of such properties are a better defence of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or viruses, and also increased tolerance of the plants to certain herbicidally active compounds. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), maize, soya beans, potatoes, cotton, tobacco, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and grapes), and particular emphasis is given to maize, soya beans, potatoes, tobacco and oilseed rape. Traits that are emphasized are in particular increased defence of the plants against insects, spiders, nematodes and slugs and snails by toxins formed in the plants, in particular those formed in the plants by the genetic material from *Bacillus thuringiensis* (for example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF and also combinations thereof) (hereinbelow referred to as "Bt plants"). Traits that are also particularly emphasized are the increased defence of the plants against fungi, bacteria and viruses by systemic acquired resistance (SAR), systemin, phytoalexins, elicitors and resistance genes and correspondingly expressed proteins and toxins.

Traits that are furthermore particularly emphasized are the increased tolerance of the plants to certain herbicidal active compounds, for example imidazolinones, sulphonylureas, glyphosate or phosphinotricin (for example the "PAT" gene). The genes which impart the desired traits in question can also be present in combination with one another in the transgenic plants. Examples of "Bt plants" which may be mentioned are maize varieties, cotton varieties, soya bean varieties and potato varieties which are sold under the trade names YIELD GARD® (for example maize, cotton, soya beans), KnockOut® (for example maize), StarLink® (for example maize), Bollgard® (cotton), Nucoton® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are maize varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosate, for example maize, cotton, soya beans), Liberty Link® (tolerance to phosphinotricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulphonylureas, for example maize). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned also include the varieties sold under the name Clearfield® (for example maize). Of course, these statements also apply to plant cultivars which have these genetic traits or genetic traits still to be developed and which will be developed and/or marketed in the future.

The plants listed can be treated in a particularly advantageous manner with the active compound mixtures according to the invention. The preferred ranges stated above for the mixtures also apply to the treatment of these plants. --

Please insert the following at page 15, between the lines 2 and 4 (compare WO 2005/009130 at page 14, lines 19-21):

-- The compounds (I) and (II) to (XIV) can be applied together, that is jointly or separately, or in succession, the sequence in the case of separate application generally not having any effect on the control results. --

Please insert the following at page 15, between the lines 7 and 9 (compare WO 2005/009130 at page 14, lines 25, through page 18, line 24):

-- Examples of suitable mixing components are the following compounds:

**Fungicides:**

2-phenylphenol; 8-hydroxyquinoline sulphate; acibenzolar-S-methyl; actinovate; aldimorph; amidoflumet; ampropylfos; ampropylfos-potassium; andoprim; anilazine; azaconazole; benalaxyl; benodanil; benomyl; benthiavalicarb-isopropyl; benzamacril; benzamacril-isobutyl; bilanafos; binapacryl; biphenyl; bitertanol; blasticidin-S; boscalid; bromuconazole; bupirimate; buthiobate; butylamine; calcium polysulphide; capsimycin; captafol; captan; carbendazim; carboxin; carvone; chinomethionat; chlobenthiazole; chlorfenazole; chloroneb; chlorothalonil; chlozolate; cis-1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol; clozylacon; cyazofamid; cyflufenamid; cymoxanil; cyproconazole; cyprodinil; cyprofuram; Dagger G; debacarb; dichlofluanid; dichlone; dichlorophen; diclomezine; dicloran; diethofencarb; difenoconazole; diflumetorim; dimethirimol; dimethomorph; dimoxystrobin; diniconazole; diniconazole-M; dinocap; diphenylamine; dipyrithione; ditalimfos; dithianon; dodine; drazoxolon; edifenphos; epoxiconazole; ethaboxam; ethirimol; etridiazole; famoxadone; fenamidone; fenapanil; fenarimol; fenbuconazole; fenfuram; fenhexamid; fenitropan; fenoxanil; fencpiclonil; fenpropidin; fenpropimorph; ferbam; fluazinam; flubenzimine; fludioxonil; flumetover; flumorph; fluoromide; fluoxastrobin; fluquinconazole; flurprimidol; flusilazole; flusulfamide; flutolanil; flutriafol; folpet; fosetyl-sodium; fuberidazole; furalaxyl; furcarbanil; furmecyclox; guazatine; hexachlorobenzene; hexaconazole; hymexazol; imazalil; imibenconazole; iminoctadine triacetate; iminoctadine tris(albesilate); iodocarb; ipconazole; iprobenfos; iprodione; iprovalicarb; irumamycin; isovaledione; kresoxim-methyl; mancozeb; maneb; meferimzone; mepanipyrim; mepronil; metalaxyl; metalaxyl-M; metconazole; methasulfocarb; methfuroxam; methyl 1-(2,3-dihydro-2,2-dimethyl-1H-inden-1-yl)-1H-imidazole-5-carboxylate; methyl 2-[[[cyclopropyl[(4-methoxyphenyl)imino]methyl]thio]methyl]-.alpha.- (methoxymethylene)benzeneacetate; methyl 2-[2-[3-(4-chlorophenyl)-1-methyl-allylideneaminooxymethyl]phenyl]-3-methoxyacrylate; metiram; metrafenone; metsulfovax; mildiomyacin; monopotassium carbonate; myclobutanil; myclozolin; nabam, N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-formylamino-2-hydroxy-benzamide; N-(6-methoxy-3-pyridinyl)cyclopropanecarboxamide; N-butyl-8-(1,1-dimethylethyl)-1-oxaspiro[4.5]decan-3-amine; natamycin; nitrothal-isopropyl; noviflumuron; nuarimol; ofurace; orysastrobin; oxadixyl; oxolinic acid; oxpoconazole; oxycarboxin;

oxyfenthiin; paclobutrazol; pefurazoate; penconazole; penthiopyrad; phosdiphen; picobenzamid; picoxystrobin; piperalin; polyoxins; polyoxorim; probenazole; prochloraz; procymidone; propamocarb; propanosine-sodium; propiconazole; propineb; proquinazid; pyraclostrobin; pyrazophos; pyrifenox; pyrimethanil; pyroxyfur; pyrrolnitrine; quinconazole; quinoxifen; quintozone; silthiofam; simeconazole; sodium tetrathiocarbonate; spiroxamine; sulphur; tecloftalam; tecnazene; tetcyclacis; tetraconazole; thiabendazole; thicyofen; thiophanate-methyl; thiram; tiadinil; tioxyimid; tolclofos-methyl; tolylfluanid; triadimefon; triazbutil; triazoxide; tricyclamide; tridemorph; trifloxystrobin; triflumizole; triforine; triticonazole; uniconazole; validamycin A; vinclozolin; zineb; ziram; zoxamide; (2S)-N-[2-[4-[[3-(4-chlorophenyl)-2-propynyl]oxy]-3-methoxyphenyl]ethyl]-3-methyl-2-[[[(methylsulphonyl)amino]-butanamide]; 1-(1-naphthalenyl)-1H-pyrrole-2,5-dione; 2,3,5,6-tetrachloro-4-(methylsulphonyl)pyridine; 2,4-dihydro-5-methoxy-2-methyl-4-[[[1-[3-(trifluoromethyl)phenyl]ethylidene]amino]oxy]methyl]phenyl]-3H-1,2,3-triazol-3-one; 2-amino-4-methyl-N-phenyl-5-thiazolecarboxamide; 2-chloro-N-(2,3-dihydro-1,1,3-trimethyl-1H-inden-4-yl)-3-pyridinecarboxamide; 3,4,5-trichloro-2,6-pyridinedicarbonitrile; 3-[(3-bromo-6-fluoro-2-methyl-1H-indol-1-yl)sulphonyl]-N,N-dimethyl-1H-1,2,4-triazole-1-sulphonamide;

and also copper salts and copper preparations, such as Bordeaux mixture; copper hydroxide; copper naphthenate; copper oxychloride; copper sulphate; cufraneb; copper(I) oxide; mancopper; oxine-copper.

### **Bactericides:**

bronopol, dichlorophen, nitrapyrin, nickel dimethyldithiocarbamate, kasugamycin, othilinone, furancarboxylic acid, oxytetracyclin, probenazole, streptomycin, tecloftalam, copper sulphate and other copper preparations.

### **Insecticides/acaricides/nematicides:**

abamectin, ABG-9008, acephate, acequinocyl, acetamiprid, acetoprole, acrinathrin, AKD-1022, AKD-3059, AKD-3088, alanycarb, aldicarb, aldoxycarb, allethrin, allethrin

1R-isomers, alpha-cypermethrin (alphamethrin), amidoflumet, aminocarb, amitraz, avermectin, AZ-60541, azadirachtin, azamethiphos, azinphos-methyl, azinphos-ethyl, azocyclotin,

Bacillus popilliae, Bacillus sphaericus, Bacillus subtilis, Bacillus thuringiensis, Bacillus thuringiensis strain EG-2348, Bacillus thuringiensis strain GC-91, Bacillus thuringiensis strain NCTC-11821, baculoviruses, Beauveria bassiana, Beauveria tenella, bendiocarb, benfuracarb, bensultap, benzoximate, beta-cyfluthrin, beta-cypermethrin, bifenazate, bifenthrin, binapacryl, bioallethrin, bioallethrin-S-cyclopentyl-isomer, bioethanomethrin, biopermethrin, bioresmethrin, bistrifluron, BMC, brofenprox, bromophos-ethyl, bromopropylate, bromfenvinfos (-methyl), BTG-504, BTG-505, bufencarb, buprofezin, butathiofos, butocarboxim, butoxycarboxim, butylpyridaben,

cadusafos, camphechlor, carbaryl, carbofuran, carbophenothion, carbosulfan, cartap, CGA-50439, chinomethionat, chlordane, chlordimeform, chloethocarb, chlorethoxyfos, chlorfenapyr, chlorfenvinphos, chlorfluazuron, chlormephos, chlorobenzilate, chloropicrin, chlorproxyfen, chlorpyrifos-methyl, chlorpyrifos (-ethyl), chlovaporthrin, chromafenozide, cis-cypermethrin, cis-resmethrin, cis-permethrin, clocythrins, cloethocarb, clofentezine, clothianidin, clothiazoben, codlemone, coumaphos, cyanofenphos, cyanophos, cycloprene, cycloprothrin, Cydia pomonella, cyfluthrin, cyhalothrin, cyhexatin, cypermethrin, cyphenothrin (1R-transisomer), cyromazine,

DDT, deltamethrin, demeton-S-methyl, demeton-S-methylsulphone, diafenthiuron, dialifos, diazinon, dichlofenthion, dichlorvos, dicofol, dicrotophos, dicyclanil, diflubenzuron, dimethoate, dimethylvinphos, dinobuton, dinocap, dinotefuran, diofenolan, disulfoton, docusat-sodium, dofenapyn, DOWCO-439,

eflusalate, emamectin, emamectin-benzoate, empenthrin (1R-isomer), endosulfan, Entomophthora spp., EPN, esfenvalerate, ethiofencarb, ethiprole, ethion, ethoprophos, etofenprox, etoxazole, etrimfos,

famphur, fenamiphos, fenazaquin, fenbutatin oxide, fenfluthrin, fenitrothion, fenobu-  
carb, fenothiocarb, fenoxacrim, fenoxycarb, fenpropathrin, fenpyrad, fenpyrithrin,  
fenpyroximate, fensulfothion, fenthion, fentrifanil, fenvalerate, fipronil, flonicamid,  
fluacrypyrim, fluazuron, flubenzimine, flubrocythrin, flucycloxuron, flucythrinate,  
flufenerim, flufenoxuron, flufenprox, flumethrin, flupyrazofos, flutenzin (flufenzine),  
fluvalinate, fonofos, formetanate, formothion, fosmethilan, fosthiazate, fubfenprox  
(fluproxyfen), furathiocarb,

gamma-HCH, gossypure, grandlure, granulosis viruses,

halfenprox, halofenozide, HCH, HCN-801, heptenophos, hexaflumuron, hexythiazox,  
hydramethylnone, hydroprene,

IKA-2002, imidacloprid, imiprothrin, indoxacarb, iodofenphos, iprobenfos, isazofos,  
isofenphos, isoprocarb, isoxathion, ivermectin,

japonilure,

kadethrin, nuclear polyhedrosis viruses, kinoprene,

lambda-cyhalothrin, lindane, lufenuron,

malathion, mecarbam, mesulfenfos, metaldehyde, metam-sodium, methacrifos,  
methamidophos, Metharhizium anisopliae, Metharhizium flavoviride, methidathion,  
methiocarb, methomyl, methoprene, methoxychlor, methoxyfenozide, metolcarb,  
metoxadiazone, mevinphos, milbemectin, milbemycin, MKI-245, MON-45700,  
monocrotophos, moxidectin, MTI-800,

naled, NC-104, NC-170, NC-184, NC-194, NC-196, niclosamide, nicotine, niten-  
pyram, nithiazine, NNI-0001, NNI-0101, NNI-0250, NNI-9768, novaluron, novi-  
flumuron,

OK-5101, OK-5201, OK-9601, OK-9602, OK-9701, OK-9802, omethoate, oxamyl, oxydemeton-methyl,

Paecilomyces fumosoroseus, parathion-methyl, parathion (-ethyl), permethrin (cis-, trans-), petroleum, PH-6045, phenothrin (1R-trans isomer), phenthoate, phorate, phosalone, phosmet, phosphamidon, phosphocarb, phoxim, piperonyl butoxide, pirimicarb, pirimiphos-methyl, pirimiphos-ethyl, prallethrin, profenofos, promecarb, propaphos, propargite, propetamphos, propoxur, prothiofos, prothoate, protrifenbute, pymetrozine, pyraclofos, pyresmethrin, pyrethrum, pyridaben, pyridalyl, pyridaphenthion, pyridathion, pyrimidifen, pyriproxyfen,

quinalphos,

resmethrin, RH-5849, ribavirin, RU-12457, RU-15525,

S-421, S-1833, salithion, sebufos, SI-0009, silafluofen, spinosad, spiroadiclofen, spiromesifen, sulfluramid, sulfotep, sulprofos, SZI-121,

tau-fluvalinate, tebufenozide, tebufenpyrad, tebupirimfos, teflubenzuron, tefluthrin, temephos, temiviphos, terbam, terbufos, tetrachlorvinphos, tetradifon, tetramethrin, tetramethrin (1R-isomer), tetrasul, theta-cypermethrin, thiacloprid, thiamethoxam, thiapronil, thiatrithos, thiocyclam hydrogen oxalate, thiodicarb, thiofanox, thiometon, thiosultap-sodium, thuringiensin, tolfenpyrad, tralocythrin, tralomethrin, transfluthrin, triarathene, triazamate, triazophos, triazuron, trichlophenidine, trichlorfon, triflumuron, trimethacarb,

vamidothion, vaniliprole, verbutin, Verticillium lecanii,

WL-108477, WL-40027,

YI-5201, YI-5301, YI-5302,

XMC, xylylcarb,

CS8726



ZA-3274, zeta-cypermethrin, zolaprofos, ZXI-8901,

the compound 3-methylphenyl propylcarbamate (tsumacide Z),

the compound 3-(5-chloro-3-pyridinyl)-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]-octane-3-carbonitrile (CAS-Reg. No. 185982-80-3) and the corresponding 3-endo-isomer (CAS-Reg. No. 185984-60-5) (cf. WO-96/37494, WO-98/25923),

and also preparations which comprise insecticidally active plant extracts, nematodes, fungi or viruses. --

Please insert new page 22 after page 21 and before the claims (compare WO 2005/009130 at pages 26-27):

-- **Example 3**

Pyricularia test (rice)/protective (II)

Rice grains (cultivar Nihonbare) were sown in plastic pots filled with sandy loam and grown in a greenhouse until they had reached the two-leaf stage.

The pots with the rice were placed in plastic beakers with the test solution and cultivated further in the greenhouse.

One or two weeks after the treatment with the test solution, the rice leaves were inoculated with Pyricularia spores ( $2.5 \times 10^5$  spores/ml)

Evaluation was carried out about one week after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Active compounds, application rates and test results are shown in the table below.

**Table 3**

Active compound	Active compound application rate in g/10 acres	Efficacy in %	Calculated efficacy in % *	Notes
<u>diclocymet</u>	20	83.2	-	a
(IX)	10	37.5	-	a
Compound	100	0	-	a
(II)				
(I)	20 } + } 100 }	100	83.2	a
+				
(IX)				
(I)	10 } + } 100 }	97.6	37.5	a
+				
(IX)				
(I)	20	75	-	b
	10	12.6	-	b
(IX)	100	27.8	-	b
(I)	20 } + } 100 }	88.1	82	b
+				
(IX)				
(I)	10 } + } 100 }	82.8	36.9	b
+				
(IX)				

a: inoculation one week after treatment

b: inoculation two weeks after treatment

\*: using Colby's formula --